

VIEWPOINT

What Other Countries Can Learn From Italy During the COVID-19 Pandemic

Stefania Boccia, MSc, PhD

Section of Hygiene, University Department of Health Sciences and Public Health, Università Cattolica del Sacro Cuore, Rome, Italy; and Department of Woman and Child Health and Public Health, Public Health Area, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy.

Walter Ricciardi, MD, MPH, MSc

Section of Hygiene, University Department of Health Sciences and Public Health, Università Cattolica del Sacro Cuore, Rome, Italy; and Department of Woman and Child Health and Public Health, Public Health Area, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy.

John P. A. Ioannidis, MD, DSc

Stanford Prevention Research Center, Department of Medicine, School of Medicine, Stanford University, Stanford, California; and Meta-Research Innovation Center at Stanford (METRICS), Stanford, California.

Corresponding

Author: John P. A. Ioannidis, MD, DSc, Stanford Prevention Research Center, 1265 Welch Rd, Medical School Office Bldg, Room X306, Stanford CA, 94305 (jioannid@stanford.edu).

In the coronavirus disease 2019 (COVID-19) pandemic, Italy has been hit very hard,¹ with 110 574 documented cases and 13 155 documented deaths related to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection as of April 1, 2020. The number of cases and deaths cannot be explained simply because of the epidemic starting in Italy earlier compared with other countries besides China. It is important to understand why death rates were so high in Italy to learn how to best prepare and how to plan for optimal actions in other countries. Some contributing factors may be immutable (eg, age structure of the population), but even these need to be laid out carefully in preparedness assessments. Some other contributing factors are potentially modifiable.

Some factors pertain to demographics and background disease in the population. Italy has the most elderly population in Europe and the second most elderly population in the world after Japan. COVID-19 has a strong age dependence for the severity of the infection and the risk of death. The median age of people infected with SARS-CoV-2 who are dying in Italy has been 80 years, and the average age of patients requiring critical care support has been 67 years. Moreover, COVID-19 morbidity and mortality is strongly dependent on the presence of concomitant serious diseases, and Italy has a high proportion of patients with history of smoking and high rates of chronic obstructive pulmonary disease and ischemic heart disease.² The corollary is that preparedness for needs of intensive care unit (ICU) beds and estimates of expected deaths should consider the age structure and chronic diseases of the population served by each health care system. Taking this adjustment into account, burden of disease may be expected to be much less in most areas in the United States, with variability across states and hospital catchment areas. For example, the proportion of the population older than 65 years is 9.5% in Alaska as compared with 19.1% in Florida and 23.1% in Italy.

A second set of factors in Italy is the increased burden of cases that presented themselves to the health care system. The proportion of people infected must have been very high in specific areas that were highly affected. In the town of Vò, all 3300 residents were tested the day the first case was detected in the third week of February, and 3% were found to be infected.³ Following aggressive testing, the epidemic was extinguished. However, elsewhere in Italy, it is likely that the prevalence of infection was several times higher in the absence of effective public health intervention. For example, it is likely that the health care system was overwhelmed in Bergamo owing to massive viral transmission during the Champions League match on

February 19, 2020 (Atalanta vs Valencia), where a third of the population of Bergamo attended and continued celebrations overnight. Italian life is famous for its socialization and frequent congregations and clustering. It is possible also that in early stages, there was not much adoption of standard hygienic measures, and instructions to stay at home proved difficult to accept, with many complaints registered with the police.⁴ Accordingly, a higher level of preparedness should be considered for areas where mass gatherings have occurred or where there is extensive social intermingling.

A third set of factors pertains to the standard capacity of the health care system and decisions made during hospital management of the presenting cases. Italy has a highly competent state-run health care system, but it has only a modest number of ICU beds and very few subintensive care beds. Overall, 5090 ICU beds (8.4 per 100 000 population) are available in Italy, and 2601 beds in coronary care units (4.3 per 100 000 population),⁵ as opposed to much higher numbers (36 ICU beds per 100 000 population) in the United States.

Given the little experience in dealing with the new virus, it is unavoidable that some strategic mistakes were made about which patients should be hospitalized. In the winter, hospitals tend to run close to full capacity, with 87% average occupancy in Italy during the flu season. Apparently, many patients with relatively modest symptoms were admitted; by the time more patients with severe cases started to arrive, there were limited reserves.

Hospital overcrowding may also explain the high infection rate of medical personnel. As of March 30, 2020, 8920 medical personnel had been found to be infected in Italy,⁶ leading to further loss of capacity for hospitals to respond. Moreover, early infection of medical personnel led to the spread of the infection to other patients within hospitals. In Lombardy, SARS-CoV-2 became largely a nosocomial infection. Nine percent of infections in Italy occurred among health care personnel.⁶ Characteristically, the first patient with COVID-19 visited the emergency department twice, thus exposing all of the personnel and patients in that area before the infection was recognized.

Italy is a decentralized country; thus, preparedness and containment may have been hampered. There was a delay from the first case detection (February 21, 2020) to the first containment decree from the government that closed the relevant villages 3 days later. The lessons relevant to other countries are the need to (1) avoid bringing patients with suspected SARS-CoV-2 infection to the hospital, except when they clearly require hospital care; (2) maintain strict hygienic procedures in the hospital environment; and (3) act swiftly in

case of exposures of medical personnel to avoid loss of personnel capacity.

Stochastic factors should also be considered. Not all of Italy, but a few cities among hundreds of cities and towns have carried most of the burden of the epidemic and have seen their hospitals crash. The Lombardy, Emilia-Romagna, and Veneto regions carry the highest numbers of infected individuals and account for 46%, 13%, and 9% of all Italian cases, respectively. The most affected provinces are Bergamo, Brescia, Milan, and Cremona, which together account for 33% of all Italian cases.⁶ There is heavy seasonality of deaths (even more so in countries with high proportions of the elderly and people who smoke, like Italy), with 25% more deaths in winter as compared with summer.⁷ Many of the excessive deaths are related to respiratory infections and are an annual occurrence. Although the infections are typically related to influenza, in 2020, SARS-CoV-2 is also a key contributor. In fact, in the 3 months prior to the outbreak, there were fewer deaths than is typical for the winter months in North Italian cities, thus leaving a larger pool of susceptible, elderly individuals.⁸ The seasonal peak of deaths varies across hospitals, and it may be difficult to predict which hospital will have the maximal burden. The corollary is that some reserves of resources, such as ventilators, should be in a stand-by allocation with the ability to assign them rapidly to hospitals that saturate their capacity.

In the absence of prevalence and incidence data, including the results of serology testing, it is difficult to predict the effects of specific major public health decisions, such as lockdowns, on the course of the COVID-19 pandemic. For example, it is not known whether

implementing a lockdown at a time when many people can infect others could lead people to spend more time in close quarters with the elderly and those who are susceptible. Similarly, it is not known whether a new epidemic wave may emerge when lockdown measures are removed. There are also unanswered questions about whether the stress and panic of a public crisis leading to major disruption and lockdown may have increased the susceptibility of elderly and frail individuals to a respiratory virus. Countries with aggressive early contact tracing and extensive laboratory testing (eg, Taiwan⁹ and South Korea) seem to offer examples of successful containment. By comparison, in Italy both contact tracing and laboratory testing were more limited, and lockdown had to be used as a last, blind measure of desperation. It is important to study the effects of the policies that are adopted first on the expected wave of patients with severe illness who will need hospitalization.

Finally, a major question that should be answered is the causal contribution of SARS-CoV-2 infection to related deaths. It is difficult to differentiate between *deaths with SARS-CoV-2 infection* and *deaths caused by SARS-CoV-2 infection* because the vast majority of patients who have died had 1 or more other major pathologies (98.8% with at least 1 comorbidity, and 48.6% having 3 or more diseases) that contributed to their death.¹⁰ Also, the lost quality-adjusted life-years of patients who died and any long-term consequences for patients who survive should be formally studied. Through this research, the relative burden of disease from COVID-19 can be better understood, and resources in overburdened health care systems during periods of crises can be better allocated.

ARTICLE INFORMATION

Published Online: April 7, 2020.

doi:10.1001/jamainternmed.2020.1447

Conflict of Interest Disclosures: None reported.

REFERENCES

1. Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *Lancet*. Published online March 13, 2020. doi:10.1016/S0140-6736(20)30627-9
2. Organization for Economic Co-Operation and Development. OECD.Stat database. Accessed March 23, 2020. <https://stats.oecd.org/>.
3. Crisanti A, Cassone A. In one Italian town, we showed mass testing could eradicate the coronavirus. *The Guardian*. March 20, 2020. Accessed April 2, 2020. <https://www.theguardian.com/commentisfree/2020/mar/20/eradicated-coronavirus-mass-testing-covid-19-italy-vo>.
4. Coronavirus, i dati dei servizi di controllo. Ministero Dell'Interno. Accessed March 23, 2020. <https://www.interno.gov.it/it/coronavirus-i-dati-dei-servizi-controllo>.
5. Epidemia COVID-19, aggiornamento nazionale: 30 marzo 2020. Task force COVID-19 del Dipartimento Malattie Infettive e Servizio di Informatica, Istituto Superiore di Sanità. Accessed April 2, 2020. https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-integrata-COVID-19_30-marzo-2020.pdf.
6. Sorveglianza integrata COVID-19: I principali dati nazionali. Istituto Superiore di Sanità. March 11, 2020. Accessed April 2, 2020. <https://www.epicentro.iss.it/coronavirus/sars-cov-2-sorveglianza-dati>.
7. Mortality monitoring in Europe. European Mortality Monitoring Project. Accessed April 3, 2020. <https://www.euromomo.eu/>.
8. Andamento della Mortalità Giornaliera (SiSMG) nelle città italiane in relazione all'epidemia di Covid-19. Centro Nazionale Prevenzione e Controllo Malattie. Accessed April 3, 2020. http://www.epiprev.it/sites/default/files/SISMG_COVID19_28032020-2.pdf
9. Wang CJ, Ng CY, Brook RH. Response to COVID-19 in Taiwan: big data analytics, new technology, and proactive testing. *JAMA*. Published online March 3, 2020. doi:10.1001/jama.2020.3151
10. Characteristics of COVID-19 patients dying in Italy. Istituto Superiore di Sanità. Accessed April 1, 2020. <https://www.epicentro.iss.it/coronavirus/sars-cov-2-decessi-italia>.